

Abyssal Current Steering of Upper Ocean Current Pathways in an Ocean Model with High Vertical Resolution

by

¹Harley E. Hurlburt, ¹E. Joseph Metzger, ¹Patrick J. Hogan,
²Charles E. Tilburg and ¹Jay F. Shriver

¹Naval Research Laboratory
Oceanography Division
Stennis Space Center, MS USA


²University of New England
Biddeford, ME USA

Ocean Sciences Meeting
Orlando, FL
3-7 Mar 2008


Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAR 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Abyssal Current Steering of Upper Ocean Current Pathways in an Ocean Model with High Vertical Resolution				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,Stennis Space Center,MS,39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Kuroshio Pathway East of Japan


Impact of topography and model resolution




1/8° 6-layer with realistic bottom topography



1/8° 6-layer flat bottom



1/4° 6-layer with realistic bottom topography



Model mean sea surface height forced by Hellerman and Rosenstein (1983, JPO) wind stress climatology
From Hurlburt et al. (1996, JGR-O; 1997, Intl WOCE Newsletter)

Bottom Current Steering of Upper Ocean Current Pathways

In a **two-layer** model, the continuity equation for layer 1 is

$$\frac{\partial h_1}{\partial t} + h_1 \nabla \cdot \vec{v}_1 + \vec{v}_1 \cdot \nabla h_1 = 0 \quad (1)$$

The **advective** term in (1) can be related to the **layer 2 velocity** by

$$\vec{v}_{1g} \cdot \nabla h_1 = \vec{v}_{2g} \cdot \nabla h_1 \quad (2)$$

$$\hat{k} \times f(\vec{v}_{1g} - \vec{v}_{2g}) = -g' \nabla h_1 \quad (3)$$

Since $|\vec{v}_1| \gg |\vec{v}_2|$ (4)

∇h_1 is a good measure of \vec{v}_1 .

From this, we see that **abyssal currents** affect the **advection** of upper layer thickness gradients and therefore the **pathways of upper layer currents**.

(Hurlburt and Thompson, 1980, JPO; Hurlburt et al., 1996, JGR-O)

Application of the 2-layer Theory for Abyssal Current Advection of Upper Ocean Current Pathways to Models with Higher Vertical Resolution

Applies when all of the following are satisfied:

- a) The flow is nearly geostrophically balanced**
- b) The barotropic and first baroclinic modes are dominant**
- c) The topography does not intrude significantly into the stratified ocean**

The interpretation in terms of surface currents applies when $|\vec{V}_{\text{near sfc}}| \gg |\vec{V}_{\text{abyssal}}|$



Notes:

- 1) The theory does not apply at low latitudes because of a) and b)**
- 2) Abyssal current advection of upper ocean current pathways is strengthened when the currents intersect at large angles, but often the end result of this advection is near barotropy**



Upper Ocean – Topographic Coupling in the Kuroshio Extension

1/12°, 20-Layer Pacific HYCOM vs. 1/8° 6-Layer NLOM

Mean SSH, RMS SSH, and mean abyssal currents




Mean abyssal currents and bottom topography




Adapted from Hurlburt et al. (2006; DAO submitted) and Hurlburt et al. (1996; JGR-O)

Global ocean depths between 200 m and 1500 m




Only 6.5% of the seafloor lies in the depth range 200-1500 m

Mean Sea Surface Temperature Around New Zealand



- Uddstrom and Oien,
JGR (1999)


Mean Currents and Sea Surface Height Simulated by (A,B) $1/16^\circ$ Linear Barotropic Model and (c) the Surface Layer from $1/8^\circ$, 6-Layer Flat Bottom NLOM




(A) QuikSCAT-corrected ECMWF ERA-40 climatological wind forcing
(B,C) Smoothed Hellerman and Rosenstein (1983) wind stress forcing

1/8°, 6-Layer NLOM Simulation of Mean Surface and Abyssal Currents East of South Island, New Zealand

Mean currents over bottom topography



Mean abyssal currents over bottom topography




— Mean subtropical front (STF)
— Mean subantarctic front (SAF)

Smoothed Hellerman and Rosenstein (1983)
(HRSM) wind forcing

From Tilburg et al. (2002; JPO)

Mean currents simulated by 1/12°, 32-layer global HYCOM in the New Zealand region overlaid on seafloor depth



Mean sea surface height in the New Zealand region

